



Primordial magnetic field constraint from 21-cm global signal

Teppei Minoda (Nagoya-U, Japan.)

with Hiroyuki Tashiro and Tomo Takahashi,
submitted to MNRAS Letters (arXiv:1812.00730)

Today's Outline

1. Introduction

- ◆ 21-cm global signal
- ◆ IGM thermal history in Dark Age
- ◆ Primordial Magnetic Fields (PMFs)

2. Theory and Methods

- ◆ Statistical property of PMFs
- ◆ MHD dissipation and heating IGM

3. Results

- ◆ IGM thermal history with PMFs
- ◆ A constraint on the PMFs



Today's Outline

1. Introduction

- ◆ 21-cm global signal
- ◆ IGM thermal history in Dark Age
- ◆ Primordial Magnetic Fields (PMFs)

2. Theory and Methods

- ◆ Statistical property of PMFs
- ◆ MHD dissipation and heating IGM

3. Results

- ◆ IGM thermal history with PMFs
- ◆ A constraint on the PMFs



21-cm line global signal

Redshift dependence of 21-cm global signal (sky averaged signal)

Roughly speaking, T_{spin} is coupled with either T_{CMB} or T_{K}

$$T_b^{21\text{cm}} \propto T_{\text{spin}} - T_{\text{CMB}} \quad T_{\text{spin}} = \frac{T_{\text{CMB}} + (y_{\text{coll}} + y_{\alpha})T_{\text{K}}}{1 + y_{\text{coll}} + y_{\alpha}}$$

(y_{coll} , y_{α} : positive coefficient)


21-cm line signal is observed as

emission when

$$T_{\text{K}} \geq T_{\text{spin}} \geq T_{\text{CMB}},$$

or absorption when

$$T_{\text{K}} \leq T_{\text{spin}} \leq T_{\text{CMB}}$$




In this work, we consider
a strong absorption around $z \sim 17$

(McQuinn & O'Leary, 2012)


IGM thermal history

Adiabatic thermal history predict absorption ($T_{\text{CMB}} > T_{\text{K}}$) at $z \sim 17$



IGM thermal history

Adiabatic thermal history predict absorption ($T_{\text{CMB}} > T_{\text{K}}$) at $z \sim 17$




Primordial Magnetic Fields

Various structures in the universe is magnetized.


- ✓ Galaxies $\sim 1\text{-}100\ \mu\text{G}$
- ✓ Galaxy Clusters $\sim 0.1\text{-}10\ \mu\text{G}$

- ✓ Voids (Intergalactic region)
 $\sim 10^{-15} - 10^{-20}\ \text{G}$

Recently, existence of magnetic fields in low-density region is suggested by gamma-ray from TeV blazars
(Ando & Kusenko 2010; Neronov & Vovk 2010)



M51 galaxy [visible & radio]
VLA/Effelsberg 20cm, HST
(Fletcher+, 2011, MNRAS, 412)



Coma cluster [radio] WSRT, 90cm
(Giovannini+, 1993, ApJ, 406)

These large-scale magnetic fields can be generated in the early universe.

(inflation, phase transition, topological defects, Harrison mechanism, ...)

= Primordial Magnetic Fields (PMFs)

(Kandus+ 2011; Subramanian 2016)

Today's Outline

1. Introduction

- ◆ 21-cm global signal
- ◆ IGM thermal history in Dark Age
- ◆ Primordial Magnetic Fields (PMFs)

2. Theory and Methods

- ◆ Statistical property of PMFs
- ◆ MHD dissipation and heating IGM

3. Results

- ◆ IGM thermal history with PMFs
- ◆ A constraint on the PMFs



stochastic PMF formalism

Assuming scale-dependent magnetic field strength

$$B_\lambda = 0 \quad (\text{for } \lambda < \lambda_{\text{cut}})$$


$$B_\lambda = B_{1 \text{ Mpc}} \left(\frac{\lambda}{1 \text{ Mpc}} \right)^{-(n_B+3)/2} \quad (\text{for } \lambda \geq \lambda_{\text{cut}})$$

scale dependence

cut-off in UV region


PMF strength
smoothed on 1Mpc

$(B_{1 \text{ Mpc}}, n_B)$ determine the
statistical property of PMFs
(and cut-off length).



*(Jedamzik+ 1998; Subramanian & Barrow 1998)

Previous constraints on PMF



PMFs as an IGM heating source

➤ Ambipolar Diffusion

- PMFs dissipate due to the friction between electrical charged particles and the neutral particles.
- Heating rate is proportional to the Lorentz force.

$$\dot{Q}_{AD} \propto |(\nabla \times \mathbf{B}) \times \mathbf{B}|^2$$

➤ Decaying Turbulence

- Kolmogorov-like turbulence cause small-scale eddies, which is thermalized due to Ohmic dissipation.
- Heating rate is proportional to the PMF energy density.

$$\dot{Q}_{DT} \propto |\mathbf{B}|^2$$

Evolutionary equations for IGM

➤ kinetic temperature

$$\frac{dT_K}{dt} = \underbrace{-2HT_K}_{\text{cosmic expansion}} + \underbrace{\frac{x_e}{1+x_e} \frac{8\rho_{\text{CMB}}\sigma_T}{3m_e c} (T_{\text{CMB}} - T_K)}_{\text{Compton heating (cooling)}} + \underbrace{\frac{\dot{Q}_{\text{AD}} + \dot{Q}_{\text{DT}}}{1.5k_B n_b}}_{\text{magnetic heating}}$$

➤ ionization fraction

$$\frac{dx_e}{dt} = \underbrace{\gamma_e n_b x_e}_{\text{collisional ionization}} + \underbrace{\left[-\alpha_e n_b x_e^2 + \beta_e (1 - x_e) \exp\left(-\frac{3E_{\text{ion}}}{4k_B T_{\text{CMB}}}\right) \right]}_{\text{photo-ionization by CMB}} \times \frac{1 + K_\alpha \Lambda n_b (1 - x_e)}{1 + K_\alpha (\Lambda + \beta_e) n_b (1 - x_e)}$$

recombination

➤ magnetic energy

$$\frac{d}{dt} \left(\frac{|B|^2}{8\pi} \right) = \underbrace{-4H \frac{|B|^2}{8\pi}}_{\text{cosmic expansion}} - \underbrace{(\dot{Q}_{\text{AD}} + \dot{Q}_{\text{DT}})}_{\text{magnetic dissipation}}$$

RECFAST code (astro-ph/9909275) + our modification

Today's Outline

1. Introduction

- ◆ 21-cm global signal
- ◆ IGM thermal history in Dark Age
- ◆ Primordial Magnetic Fields (PMFs)

2. Theory and Methods

- ◆ Statistical property of PMFs
- ◆ MHD dissipation and heating IGM

3. Results

- ◆ IGM thermal history with PMFs
- ◆ A constraint on the PMFs





IGM thermal history with PMFs

$$B_\lambda = B_{1 \text{ Mpc}} \left(\frac{\lambda}{1 \text{ Mpc}} \right)^{-(n_B+3)/2}$$

$$\frac{dT_K}{dt} = \frac{x_e}{1+x_e} \frac{8\rho_{\text{CMB}}\sigma_T}{3m_e c} (T_{\text{CMB}} - T_K) - 2HT_K$$

$$+ \frac{\dot{Q}_{\text{AD}} + \dot{Q}_{\text{DT}}}{1.5k_B n_b}$$




A constraint on PMFs

- Calculate T_K with various PMF model parameters $(B_1 \text{ Mpc}, n_B)$
- With 21-cm absorption condition as $T_K < T_{\text{CMB}}$ for $z \sim 17$, put the upper limit of PMFs.

$\Rightarrow B_1 \text{ Mpc} \lesssim 0.1 \text{ nG}$

One of the most stringent constraints on PMFs



Summary

- PMFs are possible origin of the cosmic magnetic fields
- We assume the power-law scale-dependence of PMFs
- Calculate T_K with various PMF model parameters $(B_1 \text{ Mpc}, n_B)$
- From 21-cm global absorption signal observation,
 $T_K < T_{\text{CMB}}$ for $z \sim 17$
- We derive the tightest upper bound on the PMFs as
 $B_1 \text{ Mpc} \lesssim 0.1 \text{ nG}$


Thank you for listening!




What is 21-cm line?

A radio wave due to hyperfine structure of neutral hydrogen HI


A neutral hydrogen
1s state ($n=1, l=0$)



Hyperfine structure



$$\begin{aligned}\Delta E &= 5.87 \mu\text{eV} \\ \nu &\approx 1.42 \text{ GHz} \\ \lambda &\approx 21.1 \text{ cm}\end{aligned}$$

 **21-cm line**

spin temperature $T_{\text{spin}} (\neq T_{\text{K}})$

$$\frac{n_1}{n_0} = 3 \exp\left(-\frac{\Delta E}{k_{\text{B}} T_{\text{spin}}}\right)$$

large radio telescopes (SKA, MWA, LOFAR, ...)

=> the distribution of neutral hydrogen atoms in high redshifts

=> matter density field, IGM thermal history, epoch of reionization, ...